



# Photo-period Benefit and Climate Resilient Practices on Performance and Economics of Transplanted Pigeonpea and Sunflower Intercropping in *Vertisol* of Northern Karnataka

P. Jayaraja, M.R. Umesh\*, M.Y. Ajayakumar, N. Ananda and Sreedevi S. Chavan

University of Agricultural Sciences, Raichur-584 104, India

\*E-mail: [mrumeshagri@gmail.com](mailto:mrumeshagri@gmail.com)

**Abstract:** Field experiment was conducted during *Kharif* 2019 to evaluate transplanted pigeonpea and sunflower intercropping system at different dates of sowing and fertilizers. Treatment combinations consists of two sowing time and eight intercropping system replicated thrice. Pigeonpea transplanted in second fortnight of July recorded significantly higher grain yield, equivalent yields of pigeonpea (PEY) and sunflower (SEY), over second fortnight of August. However, higher sunflower seed yield and oil content was recorded in second fortnight of August sowing. Among intercropping systems, sole pigeonpea and sunflower recorded significantly higher seed yield (2254 and 1781 kg ha<sup>-1</sup>) over sole drilled pigeonpea as well as intercropping systems. Transplanted pigeonpea + sunflower (1:2) with RDF for both crops recorded superior pigeonpea grain yield (1936 kg ha<sup>-1</sup>), PEY (2914 kg ha<sup>-1</sup>), SEY (3816 kg ha<sup>-1</sup>), gross returns (Rs. 157680 ha<sup>-1</sup>) and net returns (Rs. 108880 ha<sup>-1</sup>). Drilled pigeonpea + sunflower (1:1) considered as farmers practice recorded significantly greater sunflower seed yield (1382 kg ha<sup>-1</sup>), oil content (37.7 %) and B: C (3.49) over rest of the combinations Transplanted pigeonpea was superior over drilled sown pigeonpea.

**Keywords:** Sowing time, Transplanted pigeonpea, Drilled sown pigeonpea

Among pulses, pigeonpea [*Cajanus cajan* (L) Millsp.] holds first place in Karnataka both in area (1.48 m ha) and production (0.94 m t) with a productivity of 647 kg ha<sup>-1</sup> (Anonymous 2018). Among major production constraints, erratic and scanty rainfall that resulted low soil moisture was reducing pigeonpea productivity in *Vertisol* of N-Karnataka. Sunflower (*Helianthus annuus* L.) is grown as a source of vegetable oil; Karnataka shares 52% of the area and 40% of the production in the country. Amongst factors responsible for yield improvement in both the crops, optimum fertilizers application and timely sowing were of prime importance. Wide yield gap can be minimized by use of adequate and balanced fertilization (Umesh et al 2018, Umesh et al 2020). Crop nutrient removal far exceeds than the addition through fertilizers and manures. Balanced application of major fertilizers can increase sunflower growth and yield substantially. Long duration pigeonpea can adjust to a wide range of population and spacing. It is cultivated on marginal lands in mono/mixed cropping system without or suboptimal fertilizers under rainfed conditions. Delayed sowing shortens the time for growing period, hastens maturity and ultimately reduces the yield. The early sowing may encourage the vegetative growth which may make the crop prone to different insect pests and diseases. Thus, the need for timely planting through alternate feasible means such as transplanting is felt essential. The farmers are adopted

different row proportions in pigeonpea based intercropping rather than the recommended optimum. Hence, it was felt necessary to study the different row proportions of pigeonpea and sunflower intercropping and assess the economics of the system. Transplanting of seedlings and seed hand dibbling in pigeonpea is recently adopted techniques in the region. It has been proved that transplanting of pigeonpea would be one of the better agronomic practices to avoid delayed sowing and also maintain desired plants (Priyanka et al 2013, Praharaj et al 2015). Raising pigeonpea seedlings well in advance and transplanting in the field on receipt of good rains would help in reaping the benefits of early sowing. Keeping these facts, present investigation was carried out with an objective to evaluate feasibility of pigeonpea and sunflower intercropping in a changed sowing window and fertilizers supply in *Vertisol* of N Karnataka.

## MATERIAL AND METHODS

Field experiment was conducted in *Kharif* 2019 at Main Agricultural Research Station, Raichur. The soils of the site was medium black having soil reaction 7.74, organic carbon 0.6 % and EC of 0.21 dS m<sup>-1</sup>. The available N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O before initiation of experiment were 293.3, 39.7 and 365.9 kg ha<sup>-1</sup>. The experiment was carried out in split plot design replicated thrice. Main plots were sowing both the crops in July second fortnight and August second fortnight, sub-plot



significantly greater grain yield ( $1937 \text{ kg ha}^{-1}$ ) over drilled pigeonpea + sunflower ( $1593 \text{ kg ha}^{-1}$ ). Combined effect of sunflower intercropped with different row proportions and RDF along with time of transplanting found non-significant. Yield advantage in the early sown crop was mainly due to increased growth, yield attributes and physiological characters. It also favored by climatic conditions, temperature during growth, development and maturity stages. Efficient utilization of nutrients and moisture as well as greater light interception had resulted in higher growth and yield attributes. Similar results on yield advantage in sown crops were reported earlier worker (Reddy et al 2012, Yu 2014, Uprikar 2017). Another study at the same site by Channabasavanna et al (2015) observed that the growth and yield benefits in early sown pigeonpea due to long growing period consequently accumulated greater dry matter enhanced leaf area and phenological potential. Intercropping has reduced the yields of both pigeonpea and sunflower. Sole transplanted pigeonpea recorded higher seed and dry matter yield over intercrop with sunflower. It was mainly attributed to enhanced growth in terms of plant height, primary branches, LAI and TDMP and yield parameters viz., pods per plant, seed yield per plant, seeds per plant and 100 seed weight. Impact of intercropping on growth and yield was reported by Pal et al (2016).

**Sunflower seed yield:** The sunflower has photoperiod benefit in terms of improved yield (15.66%) either by sowing or transplanting in second fortnight of August ( $1300 \text{ kg ha}^{-1}$ ) over second fortnight of July (Table 1). Sole sunflower was out yielded ( $1781 \text{ kg ha}^{-1}$ ) over rest of the intercropped combinations. Drilled pigeonpea + sunflower (1:1) considered as farmer practice has recorded greater sunflower seed yield ( $1382 \text{ kg ha}^{-1}$ ) over sunflower as intercrop with transplanted pigeonpea (1:1) with RDF for main crop ( $831 \text{ kg ha}^{-1}$ ). Intercropping systems has reduced the sunflower yield 22.4-55.3% owing to population difference. It was compensated by contribution from pigeonpea indicated in terms of equivalent yield. Combined effect of sowing time and intercropping with row proportions and fertilizers application has no significant effect on sunflower seed yield. An uninterrupted availability of various resources viz., solar interception, soil moisture and nutrients for sole crop might have helped the crop to utilize the resources to a great extent resulting in enhanced values of varied growth parameters and yield attributes (Umesh et al 2017, Suresh et al 2019). Ahmed et al (2015) reported that sunflower yield was significantly influenced by date of sowing. The productivity of sunflower was mainly determined by the weather and soil moisture throughout its life cycle and the imposed cultural practices. Delayed sowing increased

the yields. The timely sown crop experienced moisture stress during early stage of the crop which adversely affected its growth and yield attributes. However, well distributed rainfall and adequate moisture in late sown might be responsible for enhanced sunflower yield. Additionally, sunflower is a thermo insensitive and it adjust to the late sowing but most important factor is moisture availability for their luxuriant crop growth and yield (Kaleem et al 2011, Demir 2019). Sole sunflower gave higher yields than intercropped has a result of more plant populations in the sole than intercropping (Sandeep Kumar et al 2019). Row proportion of sunflower and pigeonpea had significant effect on seed and stalk yield of sunflower. The higher seed and stalk yield of sunflower was recorded under 2:1 row proportion than 1:1 row proportion.

**Seed protein content:** Crops sown in second fortnight of July recorded significantly higher protein content (21.36 %) than second fortnight of August (20.50 %) (Table 1). Greater accumulation of nutrients in seed and better crop performance might be responsible for enhanced protein content. Sole transplanted pigeonpea recorded higher protein (22.2%) content than intercropped as well as drilled pigeonpea. Further supply of fertilizers to both the crops in transplanted pigeonpea + sunflower (1:1) also recorded greater protein content (21.32 %) over rest of the treatment combinations. Drilled pigeonpea + sunflower (1:1) and sole drilled pigeonpea was on par to each other but recorded lower than the sole transplanted and intercropped pigeonpea. Combined effect of time of sowing and intercropping system with different row proportion and fertilizer application on protein content was found non-significant. Vishwanatha (2009) reported sole pigeonpea recorded higher protein content than intercrops. In intercropping supply of balanced nutrients to the crop especially phosphorus may be due to the biochemical role of phosphorus in protein synthesis (Udikeri 2013). The protein content in seed is a function of its N concentration; therefore, higher concentration of N in seed under the superior treatments seems to be the only reason for attaining higher protein content in pigeonpea (Kumar and Paslawar 2017).

**Sunflower seed oil content:** Sunflower seed oil content was significantly greater in second fortnight of August sown (38.3%) over July (36.1%) (Table 1). Intercropping combinations were reduced the oil content over sole sunflower (39.1 %). Combined effect of time and row proportions and fertilizers application was found non-significant in respect to oil content. It was mainly due to enhanced vegetative growth and development of sunflower in August sown than July. Late-sown plants increases the seed oil might be due to the fact that the temperature increases during seed development (Allam et al 2003).

Supply of RDF to both the crops recorded higher oil content. It might be due to lower fatty acid synthesis through the pentose phosphate pathway in seeds of without fertilizer application to intercrop owing to increased competition and non-availability of required quantity of nutrients but the treatment which received RDF had better synthesis of fatty acid owing to optimum nutrient availability (Vishwanatha, 2009).

**Equivalent yields:** Pigeonpea transplanted in second fortnight of July recorded 21.1% higher PEY (2570 kg ha<sup>-1</sup>) over August (Table 1). Further it was greater in sole transplanted pigeonpea (2252 kg ha<sup>-1</sup>) than drilled pigeonpea and sunflower alone. It also improved by application of fertilizers to both the crops in transplanted pigeonpea + sunflower (1:2) (2914 kg ha<sup>-1</sup>). Application of RDF to main crop in transplanted pigeonpea + sunflower (1:1) recorded significantly lower PEY (2453 kg ha<sup>-1</sup>). Prevalence of favorable climatic weather and resources availability has resulted higher yield of both main and intercrop yield with early sowing than delayed sowing (Ravindra 2019). Murali et al (2014) also showed increased PEY with transplanting 4-5 weeks old pigeonpea seedlings than drilled sown pigeonpea as sole crop. It might be due to efficient utilization of natural resources and additional yields of component crops and their better performance in transplanted pigeonpea with higher market prices which intern resulted in higher PEY and net returns. Praharaj et al (2015) reported that significantly higher PEY over direct sown in pigeonpea and soybean intercropping and sole cropping. Intercropping system had a significant influence in obtaining higher PEY over either of

sole cropping except unfertilized control. This was due to higher seed yields of component crops owing to optimum nutrient availability (RDF to both the crops) coupled with higher price of both the crops contributed to higher PEY (Poornima 2009). Sunflower equivalent yield was 20.7% greater in crops sown in second fortnight of July (3354 kg ha<sup>-1</sup>) over August sown (Table 1). It might be due to better sunflower productivity under timely sowing than delay sowing coupled with higher market price. Intercropped treatments were recorded significantly greater SEY than the sole. Further it also enhanced by application of RDF to both the crops in transplanted pigeonpea + sunflower (1:2). It was significantly reduced under sole sunflower (1781 kg ha<sup>-1</sup>). The interaction effect between sowing time, intercropping system, row proportions and fertilizers application have significant effect on the SEY. The pigeonpea and sunflower 1:2 row proportion was recorded higher SEY as compared to other row proportions.

**Economic returns:** Crops sown/transplanted in second fortnight of July has registered higher gross returns (Rs. 136769 ha<sup>-1</sup>) net returns (Rs. 94421 ha<sup>-1</sup>) and B: C ratio (3.2) over August sown crops (Table 2). Sole transplanted pigeonpea recorded significantly higher gross returns (Rs. 123860) and net returns (Rs. 83560 ha<sup>-1</sup>) over drilled pigeonpea and sunflower. Sole Application of RDF for both crops in transplanted pigeonpea + sunflower (1:2) higher gross returns (Rs.157680 ha<sup>-1</sup>) and net returns (Rs.108880 ha<sup>-1</sup>) over 1:1 row proportions and RDF to main crop. Whereas drilled pigeonpea + sunflower (1:1) considered as farmers practice recorded higher BC ratio (3.49) over

**Table 2.** Economic returns of pigeonpea and sunflower intercropping system sown in different months and row proportions

Treatments	Cost of cultivation (Rs. ha <sup>-1</sup> )	Gross returns (Rs. ha <sup>-1</sup> )	Net returns (Rs. ha <sup>-1</sup> )	B:C ratio
Sowing /transplanting time				
July	42339	136769	94421	3.20
August	39811	114163	74361	2.85
CD (p=0.05)	-	10454	10079	0.28
Row proportions				
Transplanted pigeonpea + sunflower (1:1) RDF for main crop	43850	133285	89435	3.11
Transplanted pigeonpea + sunflower (1:1) RDF for both crops	48800	143090	94290	2.99
Transplanted pigeonpea + sunflower (1:2) RDF for main crop	43850	140430	96580	3.28
Transplanted pigeonpea + sunflower (1:2) RDF for both crops	48800	157680	108880	3.30
Drill sown pigeonpea + sunflower (1:1)	44000	142895	98895	3.49
Sole sunflower	26500	71240	44740	2.69
Sole transplanted pigeonpea	40300	123860	83560	3.07
Sole drill sown pigeonpea	32500	91245	58745	2.72
CD (p=0.05)	-	8616	7126	0.23

Market price of Pigeonpea: Rs. 55/kg, Sunflower price: Rs. 42/kg

transplanted pigeonpea + sunflower in (1:1) and 1:2. Higher grain yield which results in higher gross return same results were reported by Uprakar (2017). Higher net returns realized from intercropping was due to higher complementarity between component crops which produced higher yield. Higher returns from drilled pigeonpea + sunflower (1:1) were due to additional cost was incurred towards nursery raising and transplanting in transplanting method. Superior performance in transplanted pigeonpea treatments might be due to efficient utilization of natural resources and additional yields of component crops and their better performance in transplanted pigeonpea with higher market prices which intern resulted in higher pigeonpea equivalent yield and net returns. Similarly higher remunerative returns were observed in intercropping system over sole cropping by several workers (Vishwanatha et al 2012). Sole transplanted pigeonpea recorded higher BC ratio over sole sunflower (2.69) and drilled sown pigeonpea (2.72) (Table 2). Dibbled pigeonpea and sunflower intercropping has recorded higher BC ratio was due to higher cost of cultivation in transplanting method than of dibbling method. Similar results were reported by Priyanka et al (2013) and Ramanjaneyalu et al (2016).

### CONCLUSION

The pigeonpea either sown or transplanted was out yielded and greater economic returns. Whereas second fortnight August was benefitted to sunflower. Drill sown pigeonpea and sunflower combination in 1:1 had greater yield advantage over sole drilled pigeonpea. Yield and economic benefit were greater in second fortnight of July sown crop as compared second fortnight of August sown crop. Higher net returns was realized in transplanted pigeonpea and sunflower intercropping system with 1:2 row proportion and application of RDF based on both the crops. Transplanted pigeonpea has the greater potential and productive than drilled sown pigeonpea.

### REFERENCES

- Ahmed BN, Sultana M, Zaman J, Pau, SK, Rahman Md M, Islam MR and Majumdar F 2015. Effect of sowing dates on the yield of sunflower. *Bangladesh Agronomy Journal* **18**(1): 1-5.
- Allam AY, El-Nagar GR and Galal AH 2003. Response of two sunflower hybrids to planting dates and densities. *Acta Agronomica Hungarica* **5**(1): 25-35.
- Channabasavanna AS, Kitturmath MS and Rajkumar H 2015. Standardization of sowing date and genotypes of pigeonpea under erratic rainfall condition in northern dry zone of Karnataka. *Karnataka Journal of Agriculture Sciences* **28**(4): 604-605.
- Demir I 2019. The effects of sowing date on growth, seed yield and oil content of sunflower (*helianthus annuus l.*) cultivars under rainfed conditions. *Fresenius Environment Bulletin* **28**(9): 6849-6857
- Kaleem S, Hassan F, Ahmad M, Mahmood I, Wasaya A, Randhawa M and Khaliq P 2011. Effect of growing degree days on autumn planted sunflower. *African Journal of Biotechnology* **10**: 8840-8846.
- Kumar R and Paslawar AN 2017. Effect of conservation tillage on biomass partitioning and quality of pigeonpea based intercropping system under Vidarbha Region. *International Quarterly Journal of Life Sciences* **12**(1): 571-574.
- Murali K, Sheshadri T and Byregowda M 2014. Effect of pigeonpea transplanting on growth, yield and economics in sole and finger millet intercropping system under late sown conditions. *Journal of Food Legumes* **27**(1): 28-31.
- Pal AK, Singh R, Shukla UN and Singh S 2016. Growth and production potential of pigeonpea (*Cajanus cajan L.*) as influenced by intercropping and integrated nutrient management. *Journal of Applied and Natural Science* **8**(1): 179-183.
- Poomima DS, Shankaralingappa BC, Kalyana MKN and Savitha HR 2010. Economics of transplanted pigeonpea in sole cropping and finger millet based intercropping system. *International Journal of Agriculture Sciences* **6**(2): 501-503.
- Praharaj CS, Kumar N, Singh U, Singh SS and Singh J 2015. Transplanting in pigeonpea: A contingency measure for realizing higher productivity in Eastern Plains of India. *Journal of Food Legumes* **28**(1): 34-39.
- Priyanka K, Suresh S, Patil G, Hiremath M, Amrutha T and Sunil AK 2013. Comparative analysis of transplanted and dibbled method of Redgram cultivation in Bidar District of Karnataka. *Karnataka Journal of Agriculture Sciences* **26**: 238-242.
- Ramanjaneyulu AV, Reddy IK, Spandana BP, Neelima TL and Srinivas A 2017. Influence of pigeonpea varieties, N levels and planting methods on yield and economics under rainfed conditions. *Legume Research* **40**(5): 911-915.
- Ravindra PG 2019. *Performance of Pigeonpea Based Intercropping System under Varied Weather Condition*. Ph. D. (Agri.) Thesis, Vasantroa Naik Marathwada Krishi Vidyapeeth, Parbhani.
- Reddy MM, Padmaja B and Malathi S 2012. Evaluation of pigeonpea genotypes for delayed sowing in Telangana Region of Andhra Pradesh under Rainfed Condition. *Indian Journal of Dryland Agriculture Research Development* **27**(2): 59-62.
- Sandeep Kumar, Mandal T, Mishra GC, Barman S and Maitra S 2019. Effect of intercropping summer sunflower (*Helianthus annuus l.*) with legumes on yield attributes and productivity of crops. *International Journal of Agriculture, Environment and Biotechnology* **12**(3): 281-285.
- Singh VK and Dixit RS 1992. Effect of moisture regime and sowing date on chickpea. *Indian Journal of Agronomy* **37**: 739-743.
- Suresh SMK, Baradhan G, Elankavi S, Ramesh S and Rames N 2019. Studies on the effect of integrated nutrient management in growth attributes of sunflower in legume intercropping system. *Plant Archives* **19**(1): 1309-131.
- Udikeri 2013. *Studies on Nutrient Management in Pigeonpea and Cluster Bean Intercropping System*. M. Sc. (Agri) Thesis, UAS, Raichur.
- Umesh MR, Basavanneppa MA, Shivakumar, Mallesha, Chittapur BM, Ajayakumar MY, Desai BK and Ananda N 2020. Optimization of nitrogenous fertilizers using leaf colour chart for irrigated hybrid maize (*Zea Mays*) grown in Vertisol of sub tropical India. *Indian Journal of Agronomy* **65**: 77-82.
- Umesh MR 2002. *Assessment of Intercropping Advantage of Finger millet with Castor and Redgram under Integrated Nutrient Supply Levels*. M.Sc. thesis submitted to University of Agricultural Sciences, Bangalore.
- Umesh MR, Mallikarjun Swamy TS, Ananda N, Shanwad UK, Chittapur BM, Desai BK and Sangu Angadi 2018. Nitrogen application based on decision support tools to enhance productivity, nutrient use efficiency and quality of sweet corn (*Zea mays L.*). *Indian Journal of Agronomy* **63**(3): 331-336.
- Umesh MR, Chittapur BM and Jagadeesha N 2017. Solar radiation utilization efficiency in cereal-legume intercropping systems: A

review. *Agricultural Reviews* **38**(1): 72-75.

Uprikar AS 2017. *Performance of Pigeonpea Varieties Under Different Sowing Dates*. M.Sc. (Agri.) Thesis, Mahatma Phule Krishi Vidyapeeth, Rahuri (MH).

Vishwanatha S 2009 *Studies on Nutrient Management in Pigeonpea and Sunflower Intercropping System in 1:1 Row Proportion*. M.Sc. (Agri.), Thesis, Univ. Agric. Sci., Raichur.

Vishwanatha S, Anilkumar S N, Koppalkar B G, Pujari B T and Desai BK 2012. Effect of fertilizer management on economics and yield advantages of pigeonpea and sunflower intercropping system. *International Journal of Forestry Crop Improvement* **3**(2): 76-79.

Yu Z 2014. Two planting dates and four pigeonpea varieties at three planting densities at Taiwan. *Agriculture Science* **3**(2): 147-152.

---

Received 07 March, 2022; Accepted 22 May, 2022