



# Morphological and Chemical Profile of *Pulicaria undulata* L. (Compositae) in Iraq

Khansaa R. Al-Joboury

Iraq Natural History Research Center and Museum, University of Baghdad, Baghdad, Iraq  
E-mail: [dr.khansaa@nhm.uobaghdad.edu.iq](mailto:dr.khansaa@nhm.uobaghdad.edu.iq)

**Abstract:** The morphological features of *Pulicaria undulata* (L.) C.A. Mey. were observed. This species is a perennial plant, twiggly subshrub 30-120 cm high with a deep root system, leaves are lanceolate, semi-amplexicaul and decurrent on one side, Inflorescence in irregular cymes, peduncles 4-5 cm long, Involucre 5-7 × 3-4 mm, phyllaries numerous, linear, brown, hairy or glabrous. Kaempferol, kaempferol 3-methyl ether, 6-methoxykaempferol, quercetin, quercetin 3-methyl ether, quercetin 3,7-dimethyl were estimated using HPLC chromatography.

**Keywords:** Compositae, Chemical study, Iraq, Morphological study, *Pulicaria*

*Pulicaria undulata* (L.) C.A. Mey. (1831) (syn. *Francoeuria crispa* subsp. *crispa*, *F. crispa* var. *crispa*, *Pulicaria crispa* subsp. *crispa*, *P. crispa* var. *crispa*, *P. crispa* var. *gracillima* Maire, *P. undulata* var. *undulata*, belongs to the family compositae (asteraceae) (GBIF Secretariat, 2020). This is native in Iraq, Iran, Yemen, Afghanistan, Algeria, Benin, Chad, Djibouti, Egypt, Eritrea, Ethiopia, Gulf States, India (Dawar et al 2002). This species very important medicinal plant known as *Dethdath*, and used medicinal plants to make traditional medicines all over the world since ancient times. The branches and flowers are used to prepare powders for sneezing which works to repel insects, also in the tea of herbal (Khansaa et al 2017). This species is used for a traditional medicine which act as tonic, antispasmodic, antihypoglycemic drugs and ingredients of perfumes (Ghazanfar and Edmondson 2019) and is rich source for large number of bioactive compounds such as flavonoids which are these secondary plant metabolites and are compounds of low molecular weight. They are chemically polyphenolic and their nature offers a common structure, benzo- $\gamma$ -pyrone (Ahmed and Ibrahim 2018). These flavonoids are used by the action of antioxidants, antimicrobials, photoreceptors, optical attracts (Khalid et al 2020). This study aimed to characterize the plant through morphologic characteristics and to evaluate the flavonoids present in it.

## MATERIAL AND METHODS

Twelve plant were collected from Al-Zafaraniyah/ Baghdad- Iraq in 2020 during the flowering period in April at 3 geographical locations, the coordinate of these stations was 33°15'49.0"N 44°29'16.6"E, 33°15'44.4"N 44°29'13.1"E and

33°15'34.7"N 44°29'17.4"E. The taxonomical key was used to identify the specimens of this species for precise identification (Ghazanfar and Edmondson 2019). After being air dried in nature, the aerial parts of this species were ground in a mortar and flavonoids were isolated from *Pulicaria undulata* in pure form. Five grams of plant material separated in 100 ml of methanol using maceration (48 h). Then removed in the vacuum with the temperature at 50°C and the extracts were freeze-dried (Romanik et al 2007).

## RESULTS AND DISCUSSION

**Morphological study:** *Pulicaria undulata* is a perennial plant and twiggly subshrub 30-120 cm high with a deep root. Plant branches from the base, white-floccose or light gray in color, densely glandular, and take the form of extensive cushions. Its branches are many and may intertwine. The leaves are very long at the bottom and shorter towards the top with shape as lanceolate, semi-amplexicaul and decurrent on one side. The margin of the leaves are very clearly undulating, 2-25 × 1-5 mm, base almost auriculate (Fig. 1). Inflorescence is irregular cymes, peduncles 4-5 cm long, involucre 5-7 × 3-4 mm, phyllaries numerous, linear, brown, hairy or glabrous Ghazanfar and Edmondson (2019) also mentioned similar observation. Corolla yellow, 2.75-3 mm long, the ligule only 1 mm, and the disc florets 2.5-3.5 mm long, with glandular hairs. Cypselas 0.8 mm long, glabrous, brown, 5-ribbed, the bristles with a short narrow tuft of slightly longer hairs at the tip. Outer pappus is a ring of scales and inner pappus of bristles (short stiff hair) joined at the base into a tube, each bristle with a narrow apical tuft of slightly longer hairs. The morphology gave as clear features of this species because assemblage for species and is very useful for the

delimitation and identification. There were two types in trichomes that spread densely in its parts. The non-glandular and glandular which spread and concentrated in all parts organs: leaves, stems and flowers (Abid and Qaiser 2002) Trichome diversity were divided.

**Non-glandular trichome:** These are found in all plant parts and are simple non-glandular unicellular trichomes, different lengths ranging from short to very long about 10-860 $\mu$ m, curved all with acute apex (Fig. 2. A 1- A 3) (Krak and Mraz 2008).


**Non-glandular multicellular trichomes:** These are also found in all parts of plants, some trichome fragmented from ending with a thin flagellum, others have basal cell broader, 3-

7 cells long and the apical cell like a conical shape, others consists of two cells of different lengths, straight or curved and all with obtuse apex (Fig. 2. A 3- A7) (Al-dobaissi et al 2016).


**Stellate trichomes:** The sparse stellate hairs are on the involucre bracts (Fig. 2. A8).

**Glandular trichome:** These were multicellular divided into head and stalk (Fig. 2. B1) (Valkama et al 2003, Ciccarelli et al 2007).

**Chemical profile:** Six different flavonoids were extracted and identified: Kaempferol, kaempferol 3-methyl ether, 6-methoxykaempferol, quercetin, quercetin 3-methyl ether, quercetin 3,7-dimethyl ether (Fig. 3). The peaks for target



**Fig. 1.** Morphological characteristics of *Pulicaria undulata*: a: Whole plant, b: Root, c: the stem, d 1-3: Different lengths of leaves, e-i: Inflorescence, j: Pappus calyx



**Fig. 2.** Morphological characteristics for the trichomes of the species *Pulicaria undulata* A1-10: Non-glandular trichomes (300x), B1-5 Glandular trichomes (400x)

Kaempferol	Kaempferol 3-methyl ether	Methoxykaempferol-6
Quercetin	Quercetin 3-methyl ether	Quercetin 3,7-dimethyl ether

**Fig. 3.** Morphological form for the flavonoid compounds of the species *P. undulata*

**Table 1.** The flavonoids component with retention time and area of this species

Flavonoid compounds	Retention time (min)	Area	Concentration (mg/mL)
Kaempferol	47.82	160.55	0.26
Kaempferol 3-methyl ether	52.38	50.32	0.11
6-Methoxykaempferol	44.71	40.13	0.05
Quercetin	34.41	36.66	0.04
Quercetin 3-methyl ether	83.22	90.71	0.15
Quercetin 3,7-dimethyl ether	40.32	110.53	0.20

flavonoids eluted at retention times 47.82, 52.38, 44.71, 34.41, 83.22 and 40.32 min, respectively, through the sample extract (Table 1). Fahmied et al (2019) also made similar observation. Kaempferol recorded the highest value of 160.55 but quercetin recorded the lowest value both in retention time and area which was 36.66 and 34.41, respectively. The highest concentration recorded for kaempferol was 0.26 mg/mL. Flavonoids protect plants against different biotic and abiotic stresses, and responsible for a very important role in resistance to frost, hardness, drought resistance, also have a functional role for plant heat acclimation and freezing tolerance (Vieira et al 2003). Mierziak et al (2014) reported that the essential oil of the aerial parts for *P. gnaphalodes* has shown the presence of flavonoids, but studies on the concentrations of flavonoids are very rare in Iraq.

### CONCLUSION

The study verified the morphological features of *Pulicaria undulata* (L.) C. A. Mey. (Asteraceae) which collected from Al-Zafaraniyah/ Baghdad- Iraq. Trichomes (hairs) were divided into non-glandular and glandular. Various chemical kaempferol, kaempferol 3-methyl ether, 6-methoxykaempferol, quercetin, quercetin 3-methyl ether, quercetin 3, 7-dimethyl were estimated by HPLC chromatography. This will be helpful for medicinal use and further investigations.

### REFERENCES

Ahmed SS and Ibrahim ME 2018. Chemical investigation and antimicrobial activity of *Francoeuria crispa* (Forssk) grown wild in Egypt. *Journal of Materials and Environmental Science* **9**(1): 1-6.

Abid RD and Qaiser M 2002. Cypselae morphology of *Inula* L. (s.str.) and its allied genera (Inuleae-Compositae) from Pakistan and

Kashmir. *Pakistan Journal of Botany* **34**(3): 207-223.

Al-Dobaissi AM, Al-shimmery K and Almousawy AH 2016. Cuticular indumentum of some species from Asteraceae family in Erbil province Iraq. *Iraqi Journal of Science. Special issue-part b*: 251-262.

Ciccarelli D, Garbari F and Pagni AM 2007. Glandular hairs of the ovary: A helpful character for Asteroideae (Asteraceae) taxonomy. *Annales Botanici Fennici* **44**: 1-7.

Dawar R, Qaiser M and Perveen A 2002. Pollen morphology of *Inula* L. (s.str.) and its allied genera (Inuleae-Compositae) from Pakistan and Kashmir. *Pakistan Journal of Botany* **34**(1): 9-22.

Fahmi AA, Abdur-Rahman M, Naser AF, Hamed MA, Abd-Alla H I, Shalaby NM and Nasr MI 2019. Chemical composition and protective role of *Pulicaria undulata* (L.) C.A. Mey. subsp. *undulata* against gastric ulcer induced by ethanol in rats. *Heliyon* **5**(3): e01359.

GBIF Secretariat 2020. *Pulicaria undulata* (L.) C.A.Mey. <https://www.gbif.org/species/3095567>

Ghazanfar SA and Edmondson JR 2019. *The Flora of Iraq*, 6. 394-395. Ministry of Agriculture Republic of Iraq by Royal Botanic Gardens. Kew.

Khalid HA, Oraibi AG, Yahya HN and Al-Obaid JR 2020. Phytoremediation of Zinc Ion Using Faba Bean Plant *Vicia faba* in vitro. *Indian Journal of Ecology* **47**(4): 965-968.

Khansaa RA, Lubab GA and Sukeyna AA 2017. Morphological, anatomical and numerical taxonomy studies for some species of the fabaceae family. *Journal of Biodiversity and Environmental Sciences* **11**(5): 117-123.

Krak K and Mraz P 2008. Trichomes in tribe Lactuceae (Asteraceae)-Taxonomic implication. *Biologia* **63**(5): 616-630.

Mierziak J, Kostyn K and Kulma A 2014. Flavonoids as important molecules of plant interactions with the environment. *Molecules* **19**(10): 16240-16265.

Romanik G, Gilgenast E, Przyjazny A, Kamiński M 2007. Techniques of preparing plant material for chromatographic separation and analysis. *Journal Biochemical Biophys Methods* **70**: 253-261.

Valkama E, Salminen JP, Koricheva J and Pihlaja K 2003. Comparative analysis of leaf trichome structure and composition of epicuticular flavonoids in Finnish Birch species. *Annals of Botany* **91**: 643-655.

Vieira RF, Grayer RJ and Paton A 2003. Chemical Profiling of *Ocimum americanum* using external flavonoids. *Phytochemistry* **62**: 557.